



399PR

G-Biosciences ♦ 1-800-628-7730 ♦ 1-314-991-6034 ♦ technical@GBiosciences.com

A Geno Technology, Inc. (USA) brand name

ProteCEASE™-50, EDTA Free

Protease Inhibitor Cocktail for Large Volume

(Cat. # 786-326, 786-327)



think proteins! think G-Biosciences www.GBiosciences.com

INTRODUCTION

ProteCEASE™ -50, EDTA Free protease inhibitor cocktail contains optimized concentrations of various protease inhibitors, which provide excellent inhibition of protease activities during protein purification. ProteCEASE™ is a superior general protease inhibitor cocktail that is suitable for purification from mammalian, plant, bacteria and yeast samples. The cocktail is EDTA free and contains both irreversible and reversible protease inhibitors to inhibit serine, cysteine and other proteases. At 1X concentration, ProteCEASE™ inhibits over 90% of protease activities (e.g. Mouse Pancreas Extract 0.5mg/ml protein).

ProteCEASE™ -50, EDTA Free is designed for large volumes of extraction/ lysis buffers and is supplied as 10 vials /kit (Cat# 786-327) or 20 vials/kit (Cat# 786-327), with each vial suitable for 50ml.

ITEM(S) SUPPLIED

Description	Cat. # 786-326 For 500ml	Cat. # 786-327 For 1000ml
ProteCEASE™ -50	10 Vials	20 Vials
ProteCEASE™ Buffer	2 x 2.5ml	4 x 2.5ml

STORAGE CONDITIONS

The kit is shipped at ambient temperature. Upon arrival, store ProteCEASE™ at 4°C. If stored and used as recommended, the product is stable for 1 year.

INSTRUCTIONS FOR USE

ProteCEASE™ Buffer will freeze at 4°C. Allow all components to warm to room temperature before using. Using a clean and dry pipette tip, add 0.5ml of ProteCEASE™ Buffer to a vial of ProteCEASE™ -50 to reconstitute. Vortex briefly until completely dissolved. Close the ProteCEASE™ Buffer cap tightly. Once reconstituted, the ProteCEASE™ solution is stable for up to 1 week at 4°C.


Add 0.5ml of reconstituted ProteCEASE™ -50 directly to 50ml extraction/lysis buffer and mix it immediately for optimal protease inhibition. Once ProteCEASE™ -50 is added in the extraction/lysis buffer, use it within 20-30 minutes.

OPTIONAL: *If higher potency of protease inhibition is required, the amount of ProteCEASE™ may be doubled or sample volume reduced by half.*

NOTE: *ProteCEASE™ -50, EDTA Free is an EDTA free preparation. If metalloproteases inhibition is needed, add appropriate volume of 0.5M EDTA; e.g. 500µl 0.5M EDTA per 50 ml of sample or extraction/ lysis buffer or order our ProteCEASE™ -50, Plus EDTA kit, Cat# 786-334 or 786-335.*

RELATED PRODUCTS

Download our Protease & Phosphatase Inhibitors, Enzyme & Assays Handbook.



<http://info.gbiosciences.com/protease-phosphatase-inhibitors-enzymes-assay-handbook>

For other related products, visit our website at www.GBiosciences.com or contact us.

Last saved: 7/26/2012 CMH



www.GBiosciences.com